

Power Rule In Derivatives

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$f(x) = x^2$$

$$f'(x) = 2x$$

$$f(x) = x^2 + 3$$

$$f(x) = x^3$$


$f(x) = x^2 + 2x + 1 \Rightarrow$ Polynomial Equation

$$f(x) = x^n, \quad n \neq 0$$

if $n = 0$

$$f(x) = x^0 = 1 \Rightarrow \text{constant value}$$

$$f'(x) = 0$$



Derivative of constant = 0

$$\Rightarrow f(x) = c$$

$$f'(x) = 0$$

$$f(x) = x^n \quad \text{where } n \neq 0$$

$$f'(x) = n x^{n-1} \Rightarrow \text{Power Rule polynomial expression}$$

\Downarrow

$$\frac{\partial (f(x))}{\partial x} = n x^{n-1}$$

\Rightarrow

$$\frac{\partial (x^n)}{\partial x} = n x^{n-1}$$

$n \neq 0$

$$n=3 \quad \frac{\partial (x^3)}{\partial x} = 3 \cdot x^{3-1} = 3x^2$$

∂x

$$= 3 \times (2)^2 = 3 \times 4 = 12 //$$

$$\text{at } x=5 \quad \frac{\partial(3x^2)}{\partial x} = 3 \frac{\partial(x^2)}{\partial x} = 3 \times 2 x^{2-1} \Rightarrow 6x \Rightarrow 6 \times 5 = 30 //$$

$$\frac{\partial(1/x)}{\partial x} = \frac{\partial(x^{-1})}{\partial x} \Rightarrow -1 x^{-1-1} = -x^{-2} \Rightarrow \boxed{-\frac{1}{x^2}}$$

Assignment

$$f(x) = x^8 \Rightarrow f'(x) ?$$

$$f(x) = x^{-1} \Rightarrow f'(x) \text{ at } x = -1 //$$

④ Derivative Rules: Constant, Sum, difference And Constant Multiple

$$\frac{\partial(x^n)}{\partial x} = n x^{n-1}, \quad n \neq 0 \quad \{\text{Power Rule}\} \Rightarrow \text{polynomial}$$

$$\frac{\partial(x^0)}{\partial x} = \frac{\partial[1]}{\partial x} = 0 \Rightarrow \text{Derivative of a constant is 0.}$$

$$\frac{\partial[\overset{\text{constant}}{\downarrow} c]}{\partial x} = 0$$

$$\frac{\partial[c f(x)]}{\partial x} = c \frac{\partial(f(x))}{\partial x} = c f'(x)$$

$$\frac{\partial(3x^4)}{\partial x} = 3 \frac{\partial(x^4)}{\partial x} = 3 \times 4 x^{4-1} = 3 \times 4 x^3 = 12x^3$$

$$\text{at } x=2 \quad \frac{\partial(3x^4)}{\partial x} = 12 \times 2^3 = 12 \times 8 = 96 //$$

Assignment

$$\frac{\partial [A f(x)]}{\partial x} = \frac{\partial [2x^5]}{\partial x}$$

⊛ Sum of 2 function $f(x)$, $g(x)$

$$\frac{\partial [f(x) + g(x)]}{\partial x} \Rightarrow \frac{\partial (f(x))}{\partial x} + \frac{\partial (g(x))}{\partial x}$$

$$\begin{aligned} \frac{\partial [x^4 + x^{-2}]}{\partial x} &= \frac{\partial (x^4)}{\partial x} + \frac{\partial (x^{-2})}{\partial x} \\ &= 4x^3 + (-2x) \end{aligned}$$

$$\boxed{\frac{\partial [x^4 + x^{-2}]}{\partial x} = 4x^3 - 2x}$$


Assignment

$$\frac{\partial (x^2 + 2)}{\partial x}$$

Answer $f'(x) = \underline{\underline{2x}}$

$$\begin{aligned} \text{⊛ } \frac{\partial (4x^3 - 6x^2 + 2x + 100)}{\partial x} &\Rightarrow 12x^2 - 12x + 2 + 0 \\ &\Rightarrow \boxed{12x^2 - 12x + 2} \end{aligned}$$

Tangent of polynomials



$$\rightarrow f(x) = x^3 - 6x^2 + x - 7$$

$$y = f(1) = 1 - 6 + 1 - 7 = -11$$

$$f'(x) = \frac{\partial (x^3 - 6x^2 + x - 7)}{\partial x}$$

$$= 2x^2 - 12x + 1 - 0$$

$$f'(x) = 2x^2 - 12x + 1$$

$$f'(1) = 2 \times (1) - 12 + 1$$

$$= 2 - 12 + 1 = \boxed{-9} \Rightarrow \text{slope}$$

$$y = mx + c$$

$$y = -9(x) + c$$

$$\text{for } x = 1$$

$$-11 = -9 + c$$

$$\boxed{c = -11 + 9} \Rightarrow \boxed{c = -2}$$

$$y = mx + c$$


$$\boxed{y = -9x - 2}$$

$$m = -9 \Rightarrow \text{slope}$$

$$c = -2 \Rightarrow \text{intercept}$$

IQ Dataset

IQ	Score
100	98
90	97




{ Optimization \leftarrow



=> Chain Rule

⊕ Derivatives for Trigonometric, logarithmic and Exponential function

Trigonometric function



$$f(x) = \sin x$$

$$f'(\sin x) = \cos x$$

$$f(x) = \cos x$$

$$f'(\cos x) = -\sin x$$

logarithmic function

$$f(x) = \ln(x) \text{ then}$$

$$\boxed{f'(x) = \frac{1}{x}}$$

Exponential Function

$$f(x) = e^x$$

$$\boxed{f'(x) = e^x}$$

Constant

$$\text{if } f(x) = c$$

$$\text{then } \boxed{f'(x) = 0}$$

Power Rule

$$\text{if } f(x) = x^n$$

$$\boxed{f'(x) = nx^{n-1}}$$